

Art.-Nr.: KG1369
Ch.-B.: 134GY
Inhalt: 3 x 5 ml
Verw. bis: 2012-01

INTENDED USE

This product is intended for *in vitro* diagnostic use in the quality control of Glycerol Assays on clinical chemistry systems. The Glycerol Control is for the control of accuracy and precision.

SAFETY PRECAUTIONS AND WARNINGS

For *in vitro* diagnostic use only. Do not pipette by mouth. Exercise the normal precautions required for handling laboratory reagents.

Human source material from which this product has been derived has been tested at donor level for the Human Immunodeficiency Virus (HIV 1, HIV 2) antibody, Hepatitis B Surface Antigen (HbsAg), and Hepatitis C Virus (HCV) antibody and found to be NON-REACTIVE. FDA approved methods have been used to conduct these tests.

However, since no method can offer complete assurance as to the absence of infectious agents, this material and all patient samples should be handled as though capable of transmitting infectious diseases and disposed of accordingly.

Health and Safety Data Sheets are available on request.

STORAGE AND STABILITY

OPENED: Store refrigerated (+2 to +8°C). Reconstituted control is stable for 8 hours at 25°C or 7 days at +2 to +8°C, when kept in original container and free from contamination. Only the required amount of product should be removed. After use, any residual product should not be returned to original vial.

UNOPENED: Material stable to expiry date when stored at +2°C to +8°C.

PREPARATION FOR USE

The Glycerol control is supplied lyophilized. Carefully reconstitute each vial of Glycerol control with exactly 5 ml of distilled water at +20°C to +25°C. Close the bottle and allow to stand for 30 minutes before use. Ensure contents are completely dissolved by swirling gently. Avoid formation of foam. Do not shake.

MATERIALS PROVIDED

Glycerol Control 3 x 5 ml

MATERIALS REQUIRED BUT NOT PROVIDED

Volumetric Pipette

ASSIGNED VALUES

Values are assigned for each batch of Glycerol controls internally at BGT Laboratories Limited.

ANALYTE	TARGET	RANGE
GLYCEROL (µmol/l)	494	420 - 568
mmol/l	0.494	0.420 - 0.568
mg/dl	4.55	3.87 - 5.23

16 Sep '08